AMERICAN ACADEMY OF PEDIATRICS

Committee on Pediatric Emergency Medicine

The Use of Physical Restraint Interventions for Children and Adolescents in the Acute Care Setting

ABSTRACT. This statement defines physical restraint and provides recommendations for its use in children and adolescents.

Children and adolescents may need to be physically or chemically restrained for various procedures, because of disruptive behavior, or to prevent injury to themselves or others. The use of restraint for a child or adolescent requires clear indications, safe application, reassessment guidelines, and use only after the consideration of alternative methods. *Seclusion* refers to the involuntary confinement of a patient alone in a room, from which the patient is physically prevented from leaving, for any period of time. The use of seclusion or restraint on children and adolescents hospitalized for psychiatric disorders has been reviewed by several authors and is beyond the scope of this statement.¹⁻⁴

Restraints may be physical or chemical. Chemical restraint involves the use of psychotropic drugs or sedatives or paralytic agents. Physical restraint involves the use of physical or mechanical devices to restrain movement. Physical restraints may be cloth, leather, metal handcuffs or shackles, car seats, or seat belts. This statement is limited to the use of physical restraint in children and adolescents in the acute care or nonpsychiatric inpatient setting.

In the context of this statement, restraint differs from mechanisms that are usually and customarily used during transport or during diagnostic or surgical procedures. Common examples of these mechanisms in emergency department settings are a papoose board to aid in the control of a patient for the repair of a laceration or an arm restraint during the administration of intravenous fluids or medications.

Devices used to protect the patient, to support the patient in a specific position, or to assist in the maintenance of normal body functions are not considered restraint interventions. Examples of these types of devices are bed rails, tabletop chairs, and halter-type or soft-chest restraints.⁵

Situations that may require the short-term use of restraint in a child or adolescent include extreme, disruptive, self-injurious, or aggressive behavior as a result of drug intoxication, head injury, cerebrovas-

The recommendations in this statement do not indicate an exclusive course of treatment or serve as a standard of medical care. Variations, taking into account individual circumstances, may be appropriate.

PEDIATRICS (ISSN 0031 4005). Copyright © 1997 by the American Academy of Pediatrics.

cular hemorrhage, multiple trauma, or acute psychiatric disorder. Patients in status epilepticus may require short-term physical restraint to prevent injury to self or others until the seizure is controlled with antiepileptic agents. The use of the restraint, however, should not place a child or adolescent at risk of injury or deterioration of the medical condition.

The Joint Commission on Accreditation of Healthcare Organizations categorizes the use of restraint as a special treatment procedure requiring special justification for use. A physician's verbal or written time-limited order is obtained for each use. In addition, the standards require that hospital policy address the periodic observation of patients for whom restraint or seclusion is used.

Verbal interventions and therapeutic holding have been used for children and adolescents in psychiatric facilities to avoid the use of restraint or seclusion.^{3,4,6} *Therapeutic holding* is the physical restraint of a child by at least two people to assist the child who has lost control of behavior to regain control of strong emotions. These techniques should also be considered as options for use in the acute care setting.

RECOMMENDATIONS

Policies for the use of physical restraint of children and adolescents in the acute care or inpatient setting should include the following:

- An explanation to children why restraint is necessary, with the opportunity for children to respond to therapeutic holding when appropriate and safe.
- 2. A physician's written or verbal order specifying the type of restraint to be used and the importance of adequate restraint in relationship to its indication, with an estimate of duration. This order should be reviewed on an ongoing basis in the emergency department setting.
- 3. An immediate documented explanation to parents or family members as to why restraint is necessary.
- 4. An assessment according to hospital guidelines of those who have been restrained, assuring that the restraints are correctly applied, that skin integrity and neurovascular status remain intact, that restraints accomplish the purpose for which they were applied, and that the need for restraint continues.

Committee on Pediatric Emergency Medicine, 1995 to 1996
Joseph A. Weinberg, MD, Chairperson Barbara Barlow, MD
George L. Foltin, MD
Jerome A. Hirschfeld, MD
Dee Hodge III, MD
Jane F. Knapp, MD
William J. Lewander, MD
Karin A. McCloskey, MD
Robert A. Wiebe, MD

LIAISON REPRESENTATIVES
Jean Athey, PhD
Maternal and Child Health Bureau
Richard M. Cantor, MD
American College of Emergency Physicians
Deborah Mulligan-Smith, MD
National Association of EMS Physicians
Max L. Ramenofsky, MD
American College of Surgeons
James S. Seidel, MD
National EMSC Resource Alliance

AAP SECTION LIAISONS Patricia J. Davidson, MD Section on Anesthesiology Steven E. Krug, MD
Section on Emergency Medicine
David S. Markenson, MD
Section on Residents
Michele Moss, MD
Section on Critical Care
Dennis W. Vane, MD
Section on Surgery

REFERENCES

- Swett C Jr, Michaels AS, Cole JO. Effects of a state law on rates of restraint on a child and adolescent unit. Bull Am Acad Psychiatry Law. 1989;17:165–169
- Crespi TD. Restraint and seclusion with institutionalized adolescents. Adolescence. 1990;25:825–829
- Kalogjera IJ, Bedi A, Watson WN, Meyer AD. Impact of therapeutic management on use of seclusion and restraint with disruptive adolescent inpatients. Hosp Community Psychiatry. 1989;40:280–285
- Miller D, Walker MC, Friedman D. Use of holding technique to control the violent behavior of seriously disturbed adolescents. Hosp Community Psychiatry. 1989;40:520–524
- Special treatment procedures. In: Comprehensive Accreditation Manual for Hospitals. Oak Brook Terrace, IL: Joint Commission of the Accreditation of Healthcare Organizations; 1996:189–191
- Barlow DJ. Therapeutic holding. Effective intervention with the aggressive child. J Psychosoc Nurs Ment Health Serv. 1989;27:10–14

The Use of Physical Restraint Interventions for Children and Adolescents in the Acute Care Setting

Committee on Pediatric Emergency Medicine Pediatrics 1997;99;497 DOI: 10.1542/peds.99.3.497

Updated Information & including high resolution figures, can be found at:

Services http://pediatrics.aappublications.org/content/99/3/497

References This article cites 5 articles, 0 of which you can access for free at:

http://pediatrics.aappublications.org/content/99/3/497#BIBL

Permissions & Licensing Information about reproducing this article in parts (figures, tables) or

in its entirety can be found online at:

http://www.aappublications.org/site/misc/Permissions.xhtml

Reprints Information about ordering reprints can be found online:

http://www.aappublications.org/site/misc/reprints.xhtml





DEDICATED TO THE HEALTH OF ALL CHILDREN®



The Use of Physical Restraint Interventions for Children and Adolescents in the Acute Care Setting

Committee on Pediatric Emergency Medicine *Pediatrics* 1997;99;497 DOI: 10.1542/peds.99.3.497

The online version of this article, along with updated information and services, is located on the World Wide Web at:

http://pediatrics.aappublications.org/content/99/3/497

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 345 Park Avenue, Itasca, Illinois, 60143. Copyright © 1997 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.





DEDICATED TO THE HEALTH OF ALL CHILDREN®