

## Acetaminophen Dosing\*

NOT recommended for babies less than 2 months old.

Give every 4-6 hours as needed for fever or pain.

NOTE: Certain concentrations (mg/mL) of acetaminophen are being phased out. Please be sure to check the concentration on the bottle of acetaminophen before dosing.



Rocky Mountain  
Hospital for Children



| Weight<br>(pounds) | Weight<br>(kilogram) | Children's Liquid<br>(160 mg/5 mL) | Children's Dissolve<br>Tabs (80 mg tabs) |
|--------------------|----------------------|------------------------------------|------------------------------------------|
| Under 12 lbs       | Under 5.5            | Consult Your Physician             | —                                        |
| 12-17 lbs          | 5.5-7.9 kg           | 2.5 mL (½ teaspoon)                | —                                        |
| 18-23 lbs          | 8-10.9 kg            | 3.7 mL (¾ teaspoon)                | —                                        |
| 24-35 lbs          | 11-15.9 kg           | 5 mL (1 teaspoon)                  | 2 tabs (160 mg)                          |
| 36-47 lbs          | 16-20.9 kg           | 7.5 mL (1 ½ teaspoons)             | 3 tabs (240 mg)                          |
| 48-59 lbs          | 21-26.5 kg           | 10 mL (2 teaspoons)                | 4 tabs (320 mg)                          |
| 60-71 lbs          | 26.6-32.2 kg         | 12.5 mL (2 ½ teaspoons)            | 5 tabs (400 mg)                          |

\* For Example, Tylenol

## Ibuprofen Dosing\*

For infants and children 6 months of age and older.

Give every 6-8 hours as needed for fever or pain.


| Weight<br>(pounds) | Weight<br>(kilogram) | Children's Liquid<br>(100 mg/5 mL) | Junior tablets<br>(100 mg) | Adult tablets<br>(200 mg) |
|--------------------|----------------------|------------------------------------|----------------------------|---------------------------|
| Under 12 lbs       | Under 5.5            | Consult Your Physician             | —                          | —                         |
| 12-21 lbs          | 5.5-9.9 kg           | 2.5 mL (½ teaspoon)                | —                          | —                         |
| 22-33 lbs          | 10-14.9 kg           | 5 mL (1 teaspoon)                  | 1 tablet (100 mg)          | —                         |
| 34-43 lbs          | 15-19.9 kg           | 7.5 mL (1 ½ teaspoons)             | 1 tablet (100 mg)          | —                         |
| 44-55 lbs          | 20-24.9 kg           | 10 mL (2 teaspoons)                | 2 tablets (200 mg)         | 1 tablet (200 mg)         |
| 56-66 lbs          | 25-29.9 kg           | 12.5 mL (2 ½ teaspoons)            | 2 tablets (200 mg)         | 1 tablet (200 mg)         |
| 67-88 lbs          | 30-39.9 kg           | 15 mL (3 teaspoons)                | 3 tablets (300 mg)         | —                         |
| 89+ lbs            | 40+ kg               | —                                  | 4 tablets (400 mg)         | 2 tablets (400 mg)        |

\* For Example, Motrin and Advil

Where the flu is foiled.

RMHC033 (07/22)

## Rocky Mountain Hospital for Children ER Locations



**1** ROCKY MOUNTAIN HOSPITAL FOR CHILDREN  
at Presbyterian/St. Luke's Medical Center  
Full Service Pediatric Hospital | 24/7 Peds ER  
20th Ave & High Street | 720-754-1000

**2** Rocky Mountain Hospital for Children  
at Swedish Medical Center  
Level I Trauma Center | Burn Program  
NICU | Pediatric Specialty ICU

### Pediatric Services Available at Affiliated Hospitals

- 3** North Suburban Medical Center  
NICU
- 4** Rose Medical Center  
NICU
- 5** The Medical Center of Aurora  
Child/Adolescent Behavioral Health  
NICU
- 6** Sky Ridge Medical Center  
NICU | Pediatric Unit

WeAreAmazingTogether.com

### Dosis de Acetaminofén\*

NO se recomienda para bebés menores de 2 meses de edad.  
Dar medicamento cada 4-6 horas para aliviar síntomas de fiebre o dolor.

NOTA: En algunas concentraciones (mg/mL) de acetaminofén las fases se separan. Asegúrese de verificar la concentración en el frasco de acetaminofén antes de administrarla.



Rocky Mountain  
Hospital for Children



| Peso (libras)   | Peso (kilos)    | Jarabe líquido para niños (160 mg/5 mL) | Tabletas masticables (tableta de 80 mg) |
|-----------------|-----------------|-----------------------------------------|-----------------------------------------|
| Menos de 12 lbs | Menos de 5.5 kg | Consulte a Su Pediatra                  | —                                       |
| 12-17 libras    | 5.5-7.9 kilos   | 2.5 mL (½ cucharita)                    | —                                       |
| 18-23 libras    | 8-10.9 kilos    | 3.7 mL (¾ cucharita)                    | —                                       |
| 24-35 libras    | 11-15.9 kilos   | 5 mL (1 cucharita)                      | 2 tabletas (160 mg)                     |
| 36-47 libras    | 16-20.9 kilos   | 7.5 mL (1 ½ cucharitas)                 | 3 tabletas (240 mg)                     |
| 48-59 libras    | 21-26.5 kilos   | 10 mL (2 cucharitas)                    | 4 tabletas (320 mg)                     |
| 60-71 libras    | 26.6-32.2 kilos | 12.5 mL (2 ½ cucharitas)                | 5 tabletas (400 mg)                     |

\* Por ejemplo, Tylenol

### Dosis de Ibuprofeno\*

Para niños mayores de 6 meses.  
Dar medicamento cada 6-8 horas para aliviar síntomas de fiebre o dolor.


| Peso (libras)   | Peso (kilos)    | Jarabe líquido para niños (100 mg/5 mL) | Tabletas para niños (100 mg) | Pastillas para adultos (200 mg) |
|-----------------|-----------------|-----------------------------------------|------------------------------|---------------------------------|
| Menos de 12 lbs | Menos de 5.5 kg | Consulte a Su Pediatra                  | —                            | —                               |
| 12-21 libras    | 5.5-9.9 kilos   | 2.5 mL (½ cucharita)                    | —                            | —                               |
| 22-33 libras    | 10-14.9 kilos   | 5 mL (1 cucharita)                      | 1 tablet (100 mg)            | —                               |
| 34-43 libras    | 15-19.9 kilos   | 7.5 mL (1 ½ cucharitas)                 | 1 tablet (100 mg)            | —                               |
| 44-55 libras    | 20-24.9 kilos   | 10 mL (2 cucharitas)                    | 2 tablets (200 mg)           | 1 pastilla (200 mg)             |
| 56-66 libras    | 25-29.9 kilos   | 12.5 mL (2 ½ cucharitas)                | 2 tablets (200 mg)           | 1 pastilla (200 mg)             |
| 67-88 libras    | 30-39.9 kilos   | 15 mL (3 cucharitas)                    | 3 tablets (300 mg)           | —                               |
| 89+ libras      | 40+ kilos       | —                                       | 4 tablets (400 mg)           | 2 pastillas (400 mg)            |

\* Por ejemplo, Motrin y Advil

Where the flu is foiled.

RMHC033 (07/22)

### Rocky Mountain Hospital for Children ER Locations



- 1** **ROCKY MOUNTAIN HOSPITAL FOR CHILDREN at Presbyterian/St. Luke's Medical Center**  
Full Service Pediatric Hospital | 24/7 Peds ER  
20th Ave & High Street | 720-754-1000
- 2** **Rocky Mountain Hospital for Children at Swedish Medical Center**  
Level I Trauma Center | Burn Program  
NICU | Pediatric Specialty ICU

#### Pediatric Services Available at Affiliated Hospitals

- 3** **North Suburban Medical Center**  
NICU
- 4** **Rose Medical Center**  
NICU
- 5** **The Medical Center of Aurora**  
Child/Adolescent Behavioral Health  
NICU
- 6** **Sky Ridge Medical Center**  
NICU | Pediatric Unit

WeAreAmazingTogether.com